

A LIST OF THE BIRDS COLLECTED ON THE
ISLAND OF TOBAGO, WEST INDIES, BY W. W.
BROWN, JR., DURING APRIL
AND MAY, 1892.

BY CHARLES B. CORY.

THE collection made by Mr. Brown contained over seven hundred bird skins and a few mammals, including two specimens of *Sciurus griseogenys* Gray. The birds are especially interesting, several of them being new to the island.

- | | |
|--|--|
| <i>Sterna elegans</i> Gamb. | <i>Pachyrhamphus albo-griseus</i> Schl. |
| <i>Ardea virescens</i> Linn. | <i>Pachyrhamphus niger</i> Spix. |
| <i>Ionornis martinica</i> (Linn.). | <i>Chiroxiphia pareola</i> (Linn.). |
| <i>Ortalis ruficauda</i> Fard. | <i>Tyrannus melancholicus</i> Vieill. |
| <i>Columba rufina</i> (Temm.). | <i>Myiarchus tyrannulus</i> (Müll.). |
| <i>Engyptila verreauxi</i> (Bp.). | <i>Elainea pagana</i> (Licht.). |
| <i>Columbigallina rufipennis</i> (Bp.). | <i>Ostinops decumanus</i> (Pall.). |
| <i>Geotrygon linearis</i> (Knip et Prev.). | <i>Molothrus atronitens</i> (Cab.). |
| <i>Buteo latissimus</i> Wils. | <i>Euetheia bicolor</i> (Linn.). |
| <i>Spizaetus manduyti</i> Daud. | <i>Spermophila lineata</i> (Gm.). |
| <i>Amazona amazonica</i> (Linn.). | <i>Volatinia splendens</i> (Vieill.). |
| <i>Crotophaga ani</i> Linn. | <i>Tanagra sclateri</i> Berlepsch. |
| <i>Dendrobates kirki</i> (Malh.). | <i>Tachyphonus melaleucus</i> Sparrm. |
| <i>Melanerpes terricolor</i> Berlepsch. | <i>Vireo chivi</i> (Vieill.). |
| <i>Chloronerpes rubiginosus</i> (Sw.). | <i>Hylophilus insularis</i> Sclater. |
| <i>Trogon collaris</i> Vieill. | <i>Progne dominicensis</i> (Gmel.). |
| <i>Momotus swainsoni</i> Sclater. | <i>Cæreba luteola</i> (Cab.). |
| <i>Ceryle americana</i> Gm. | <i>Arbelorhina cyanea</i> (Linn.). |
| <i>Galbula ruficauda</i> (Cuv.). | <i>Troglodytes tobagensis</i> Lawr. |
| <i>Chrysolampis moschitus</i> (Linn.). | <i>Thryothorus rutilus</i> (Vieill.). |
| <i>Florisuga mellivora</i> (Linn.). | <i>Mimus gilvus</i> Vieill.—In a series of |
| <i>Amazilia erythronota</i> (Less.). | <i>Mimus</i> from Tobago all the speci- |
| <i>Campylopterus ensipennis</i> (Swain.) | mens are paler and more ashy in |
| <i>Formicivora intermedia</i> Licht. | color on the upper parts than birds |
| <i>Dysithamnus semicinereus</i> Sclater. | from Grenada and St. Vincent. |
| <i>Thamnophilus doliatus</i> (Linn.). | <i>Merula xanthosceles</i> (Fard.). |
| <i>Dendroornis sussurans</i> Fard. | <i>Merula gymnophthalma</i> (Caban.). |